

Always ask your healthcare professional which product is right for your child.

Your child's weight and age

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving MOTRIN[®]. A healthcare professional should be consulted for dosing in children under the age of six months.

	Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
	Age	0-5 mos	6-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs
 <p>Concentrated MOTRIN[®] Infants' Drops Berry & Berry dye-free flavors Active Ingredient: Ibuprofen 50mg (in each 1.25mL) Use only as directed.</p>		—	1.25mL	1.875mL	—	—	—	—	—
 <p>Children's MOTRIN[®] Suspension Berry, Grape, Tropical Punch, Bubblegum & Berry dye-free flavors Active Ingredient: Ibuprofen 100mg (in each 5mL or 1 tsp) Use only as directed.</p>		—	—	—	1 tsp or 5mL	1½ tsp or 7.5mL	2 tsp or 10mL	2½ tsp or 12.5mL	3 tsp or 15mL
 <p>MOTRIN[®] Junior Strength Easy-to-Swallow Caplets Active Ingredient: Ibuprofen 100mg (in each caplet) Use only as directed.</p>		—	—	—	—	—	2 caplets	2½ caplets	3 caplets
 <p>MOTRIN[®] Junior Strength Chewable Tablets Orange & Grape flavors Active Ingredient: Ibuprofen 100mg (in each tablet) Use only as directed.</p>		—	—	—	1 tablet	1½ tablets	2 tablets	2½ tablets	3 tablets
Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours.									

Important Instructions for Proper Use

- Read and follow the label instructions on all MOTRIN[®] products
 - Take every 6-8 hours as needed
 - Do not exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or non-prescription)
 - MOTRIN[®] contains ibuprofen
- Keep all medicines out of the reach of children

- Use only the dosing device that comes with a specific product
 - Household items such as measuring spoons are less accurate
- Do NOT administer adult medicines to children
- Concentrated MOTRIN[®] Infants' Drops are more concentrated than Children's MOTRIN[®] Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dosing device
- Children's MOTRIN[®] Liquids are less concentrated than Concentrated MOTRIN[®] Infants' Drops. The Children's MOTRIN[®] Liquids have been specifically designed for use with the enclosed measuring cup
- The OTC dosing: the recommended dose is 7.5mg/kg every 6-8 hours. The recommended daily dose is 30mg/kg
- If you have any questions, contact your healthcare professional or call 1-877-895-3665

Dosing for Infants and Children

From Your Healthcare Professional

Infants' **TYLENOL**

Children's **TYLENOL**

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise use age.

mL = milliliter
tsp = teaspoon



Infants' TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL)

Available in:
 Grape
 Cherry

Use only as directed.



Children's TYLENOL® Oral Suspension

Active Ingredient:
Acetaminophen 160 mg
(in each 5 mL or 1 tsp)

Available in:
 Grape
 Cherry

Use only as directed.

WEIGHT	AGE	Infants' TYLENOL®	Children's TYLENOL®
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)